

Some Remarks on Turán's Inequality

S. P. ZHOU

*Department of Mathematics, Statistics and Computing Science,
Dalhousie University, Halifax, NS Canada B3H 3J5*

Communicated by T. J. Rivlin

Received July 13, 1990; revised January 16, 1991

We examine how large the L^p norm on $[-1, 1]$ of the derivative of a real algebraic polynomial of degree at most n with zeros only in $[-1, 1]$ can be if the L^q norm of the polynomial is 1. © 1992 Academic Press, Inc.

Let H_n be the class of real algebraic polynomials of degree n , whose zeros all lie in the interval $[-1, 1]$. Let R_n be the class of real trigonometric polynomials of degree n with only real roots. Let

$$\|f\|_{C[a,b]} = \|f\|_{L^\infty[a,b]} = \max_{a \leq x \leq b} |f(x)|,$$

$$\|f\|_{L^p[a,b]} = \left(\int_a^b |f(x)|^p dx \right)^{1/p} \quad \text{for } 1 \leq p < \infty.$$

In 1939, P. Turán [4] proved that for $f \in H_n$,

$$\|f'\|_{C[-1,1]} \geq Cn^{1/2} \|f\|_{C[-1,1]},$$

where here and throughout the whole paper, C always indicates some positive absolute constant. Later, J. Erőd [3] improved the constant above. In 1976, A. K. Varma [5] established the corresponding inequality in L^2 space. The general form for $1 \leq p \leq \infty$ of the inequality is due to the author [7]:

THEOREM 1. *If $f \in H_n$, then for $1 \leq p \leq \infty$,*

$$\|f'\|_{L^p[-1,1]} \geq Cn^{1/2} \|f\|_{L^p[-1,1]}.$$

In 1983, A. K. Varma [6] obtained the best possible constant in this inequality when $p = 2$. In 1986, V. F. Babenko and S. A. Pichugov [1, 2]

found the best possible constants in this inequality and in the corresponding trigonometric case when $p = \infty$.

In [8] we considered similar results in L^p for $0 < p < 1$.

The object of this paper is to establish the following more general result.

THEOREM 2. *If $f \in H_n$, then for $1 \leq p \leq q \leq \infty$,*

$$\|f'\|_{L^p[-1,1]} \geq C n^{1/2 - 1/(2p) + 1/(2q)} \|f\|_{L^q[-1,1]}. \quad (*)$$

Proof. Denote by $-1 \leq x_1 < x_2 < \dots < x_k \leq 1$ all the distinct zeros of $f \in H_n$, and the order of x_i by m_i . Write the maximum point of $|f(x)|$ between (x_i, x_{i+1}) as α_i . If $x_1 > -1$ or $x_k < 1$, set $\alpha_0 = -1$ or $\alpha_k = 1$. Also, let

$$m(x) = \sum_{i=1}^k \frac{m_i}{x - x_i}.$$

We divide the proof into several cases.

Case 1. $\alpha_j - x_j \leq n^{-1/2}$ for some j . Then applying Hölder's inequality we have

$$\int_{x_j}^{\alpha_j} |f'(x)|^p dx \geq n^{(p-1)/2} |f(\alpha_j)|^p \geq n^{(p-1)/2} (\alpha_j - x_j)^{-p/q} \left(\int_{x_j}^{\alpha_j} |f(x)|^q dx \right)^{p/q},$$

that is,

$$\int_{x_j}^{\alpha_j} |f'(x)|^p dx \geq n^{(pq-q+p)/(2q)} \left(\int_{x_j}^{\alpha_j} |f(x)|^q dx \right)^{p/q}.$$

If $\alpha_j - x_j \geq n^{-1/2}$ for some j , put $d_j = [(\alpha_j - x_j)n^{1/2}]$, where $[x]$ denotes the greatest integer not exceeding x .

Case 2.1. $\alpha_j - x_j > n^{-1/2}$ for some j . Then as above,

$$\int_{x_j}^{\alpha_j - d_j/n^{1/2}} |f'(x)|^p dx \geq n^{(pq-q+p)/(2q)} \left(\int_{x_j}^{\alpha_j - d_j/n^{1/2}} |f(x)|^q dx \right)^{p/q}.$$

Case 2.2. $\alpha_j - x_j > n^{-1/2}$ for some j , and $|f(\alpha_j - tn^{-1/2})| \geq 2|f(\alpha_j - (t+1)n^{-1/2})|$ for some $t = 0, 1, \dots, d_j - 1$. Using Hölder's inequality again, we have

$$\int_{\alpha_j - (t+1)/n^{1/2}}^{\alpha_j - t/n^{1/2}} |f'(x)|^p dx \geq n^{(p-1)/2} |f(\alpha_j - tn^{-1/2}) - f(\alpha_j - (t+1)n^{-1/2})|^p,$$

and since

$$|f(\alpha_j - tn^{-1/2})| \geq 2 |f(\alpha_j - (t+1)n^{-1/2})|,$$

we get

$$\int_{\alpha_j - (t+1)/n^{1/2}}^{\alpha_j - t/n^{1/2}} |f'(x)|^p dx \geq 2^{-p} n^{(pq-q+p)/(2q)} \left(\int_{\alpha_j - (t+1)/n^{1/2}}^{\alpha_j - t/n^{1/2}} |f(x)|^q dx \right)^{p/q}.$$

Case 2.3. $\alpha_j - x_j > n^{-1/2}$ for some j , and $|f(\alpha_j - rn^{-1/2})| < 2 |f(\alpha_j - (r+1)n^{-1/2})|$ for some $r=0, 1, \dots, d_j-1$. Applying the mean value theorem to the function $f'(x)/f(x)$ on the interval $(x_j, \alpha_j]$ and noting that $f'(\alpha_j)=0$, we have for $x \in (x_j, \alpha_j - \frac{1}{2}n^{-1/2})$, $j \neq k$,

$$m(x) = \sum_{i=1}^k \frac{m_i}{x - x_i} = (\alpha_j - x) \sum_{i=1}^k \frac{m_i}{(\xi - x_i)^2} \geq \frac{n^{1/2}}{8}, \quad \xi \in (x, \alpha_j);$$

a similar calculation leads to

$$m(x) \geq \sum_{i=1}^k \frac{m_i}{x - x_i} - \sum_{i=1}^k \frac{m_i}{\alpha_k - x_i} \geq \frac{n^{1/2}}{8}$$

for $x \in (x_k, \alpha_k - \frac{1}{2}n^{-1/2})$. So

$$\begin{aligned} \int_{\alpha_j - (r+1)/n^{1/2}}^{\alpha_j - r/n^{1/2}} |f'(x)|^p dx &\geq \int_{\alpha_j - (r+1)/n^{1/2}}^{\alpha_j - (2r+1)/(2n^{1/2})} |f(x)|^p |m(x)|^p dx \\ &\geq 2^{-3p-1} n^{(p-1)/2} |f(\alpha_j - (r+1)n^{-1/2})|^p \\ &\geq 2^{-4p-1} n^{(p-1)/2} |f(\alpha_j - rn^{-1/2})|^p; \end{aligned}$$

hence

$$\int_{\alpha_j - (r+1)/n^{1/2}}^{\alpha_j - r/n^{1/2}} |f'(x)|^p dx \geq 2^{-4p-1} n^{(pq-q+p)/(2q)} \left(\int_{\alpha_j - (r+1)/n^{1/2}}^{\alpha_j - r/n^{1/2}} |f(x)|^q dx \right)^{p/q}.$$

Now Cases 2.1–2.3 prove that

$$\int_{x_j}^{\alpha_j} |f'(x)|^p dx \geq 2^{-4p-1} n^{(pq-q+p)/(2q)} \left(\int_{x_j}^{\alpha_j} |f(x)|^q dx \right)^{p/q}$$

for $\alpha_j - x_j > n^{-1/2}$, while Case 1 proves the inequality if $\alpha_j - x_j \leq n^{-1/2}$. The same inequality clearly holds if $\int_{x_j}^{\alpha_j}$ is replaced with $\int_{x_j}^{x_j+1}$. Summing over all j , we get the claimed inequality. ■

With $f(x) = (1-x^2)^{[n/2]}$, we see that the order $n^{1/2-1/(2p)+1/(2q)}$ in (*) cannot be improved.

The proofs for the following theorems are similar.

THEOREM 3. Let $1 \leq p \leq q \leq \infty$, and $f(x)$ be an n th degree algebraic polynomial, which has only real roots. If at most k roots of $f(x)$ lie outside the interval $[-1, 1]$, then

$$\|f'\|_{L^p[-1,1]} \geq C_k n^{1/2 - 1/(2p) + 1/(2q)} \|f\|_{L^q[-1,1]},$$

where C_k is a positive constant depending only upon k .

THEOREM 4. Let $1 \leq p \leq q \leq \infty$. If $f \in R_n$, then

$$\|f'\|_{L^p[0,2\pi]} \geq C n^{1/2 - 1/(2p) + 1/(2q)} \|f\|_{L^q[0,2\pi]}.$$

ACKNOWLEDGMENTS

The author thanks Dr. Peter Borwein for his valuable discussions and comments. Many thanks are also due to the referees for their helpful comments.

REFERENCES

1. V. F. BABENKO AND S. A. PICHUGOV, Accurate inequality for the derivatives of trigonometric polynomials, which have only real zeros, *Math. Notes* **39** (1986), 179–182.
2. V. F. BABENKO AND S. A. PICHUGOV, Inequality for the derivatives of polynomials with real zeros, *Ukrain. Math. J.* **38** (1986), 347–351.
3. J. ERŐD, Bizonyos polinomok maximumairól, *Math. Fiz. Lapok* **46** (1939), 58–82.
4. P. TURÁN, Über die Ableitung von Polynomen, *Compositio Math.* **7** (1939), 89–95.
5. A. K. VARMA, An analogue of some inequalities of P. Turán concerning algebraic polynomials satisfying certain conditions, *Proc. Amer. Math. Soc.* **55** (1976), 305–309.
6. A. K. VARMA, Some inequalities of algebraic polynomials having all zeros inside $[-1, 1]$, *Proc. Amer. Math. Soc.* **88** (1983), 227–233.
7. S. P. ZHOU, On Turán's inequality in L^p norm (Chinese), *J. Hangzhou Univ.* **11** (1984), 28–33. MR 85j:26025.
8. S. P. ZHOU, An extension of the Turán inequality in L^p for $0 < p < 1$, *J. Math. Res. Exposition* **6**, No. 2 (1986), 27–30. MR 89c:41011.